

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 031 627 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
30.08.2000 Bulletin 2000/35

(21) Application number: 00102289.6

(22) Date of filing: 17.02.2000

(51) Int. Cl.⁷: **C12N 15/53**, C12N 9/02,
C12N 1/21, C12N 5/10,
C12P 23/00, C07K 16/40,
C12Q 1/68, G01N 33/573
// (C12N1/21, C12R1:19)

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: 22.02.1999 EP 99103382

(71) Applicant:
F. HOFFMANN-LA ROCHE AG
4070 Basel (CH)

(72) Inventors:
• **Bachmann, Heinrich**
4451 Wintersingen (CH)
• **Brugger, Roland**
79106 Freiburg (DE)
• **Friedlein, Arno Martin**
79189 Bad Krozingen (DE)

- **Wirtz, Gabriele Margarethe**
42349 Wuppertal (DE)
- **Woggon, Wolf-Dietrich**
4102 Binningen (CH)
- **Wyss, Adrian**
4313 Möhlin (CH)
- **Wyss, Markus**
4419 Liestal (CH)

(74) Representative:
Keller, Günter, Dr. et al
Lederer, Keller & Riederer
Patentanwälte
Prinzregentenstrasse 16
80538 München (DE)

Remarks:

The applicant has subsequently filed a sequence listing and declared, that it includes no new matter.

(54) **Beta-carotene 15,15'-dioxygenase**

(57) Nucleic acid sequences coding for a protein having β,β -carotene 15,15'-dioxygenase activity and their uses in diagnostics, the synthesis of vitamin A and methods for the introduction of the β,β -carotene 15,15'-dioxygenase cDNA into host cells are disclosed.


EP 1 031 627 A1

Description

[0001] The present invention concerns the cloning of β,β -carotene 15,15'-dioxygenase (EC 1.13.11.21), the enzyme responsible for the cleavage of β -carotene leading to vitamin A. The term vitamin A as defined in the present invention comprises a class of compounds including retinal, retinol, 3-dehydroretinol, retinoic acid, the isomers from these compounds as well as retinylesters. Proteins having β,β -carotene 15,15'-dioxygenase activity and nucleic acid sequences coding therefore can be used in different fields including but not limited to diagnostics, the technical production of vitamin A, the generation of transgenic plants in order to produce vitamin A in fruits and vegetables, or gene therapy.



[0002] Vitamin A is essential for man and animal and is largely formed in most organisms from its precursor carotenoids which, by themselves, can only be formed in plants, in photo-synthetic active and some other microorganisms. Man and most animals (in particular herbivores and omnivores) are able to convert such carotenoids, also called provitamins A, enzymatically into vitamin A. The most important enzyme for this process is the β,β -carotene 15,15'-dioxygenase (EC 1.13.11.21). The enzyme is located in the cytosol and forms retinal from β -carotene, as the principal substrate, in presence of oxygen according to scheme 1:

Scheme I:



[0003] The enzyme β,β -carotene 15,15'-dioxygenase is characterized by generating 2 mols retinal from 1 mol of β -carotene by central cleavage. But the enzyme is also able to convert a wide range of carotenoids into vitamin A-active compounds, as shown in scheme II:

Scheme II:



[0004] Highest known enzymatic activity is found in the intestine of herbivores, especially in duodenum. In other tissues like liver, lung, kidney and brain β,β -carotene 15,15'-dioxygenase is also detectable. Starting in 1955, many attempts have been undertaken to purify and characterize the enzyme by biochemical methods [Goodman (1965 and 1966), Fidge (1969), Laksmanan (1972), Sharma (1977) and Devery & Milborrow (British Journal of Nutrition (1994) 72, p. 397-414]. However, none of these attempts has been successful. Specific activities of 600 pmol retinal formed/mg protein per hour have not been surpassed.

[0005] In the course of the present invention it was possible to purify the chicken enzyme to such a degree that a partial amino acid sequence could be obtained. The enzyme was enriched 226-fold, yielding a specific activity of 2500 pmol/h/mg. On a polyacrylamid gel of fractions from the final gel filtration run 15 bands were visible after Coomassie

blue staining. Two bands correlated with the enzymatic activity profile. With the first protein Edman sequencing and with the second MS spectroscopy was performed. Tryptic ingel digestion and subsequent microbore RP-HPLC peptide mapping in combination with MALDI-TOF MS and automated Edman degradation of this latter protein revealed 2 peptides of 11 and 18 amino acids. From this sequence information degenerate PCR primers were designed and synthesized.

[0006] With a PCR protocol a 51 bp (base pair) fragment was amplified within the longer peptide. From this sequence a homologous primer was synthesized and used in a second RT-PCR (reverse transcriptase - PCR) to amplify a 597 bp fragment.

[0007] This cDNA fragment was radioactively labeled and used for the screening of two positive pools from a chicken expression library in order to isolate the full length cDNA coding for β,β -carotene 15,15'-dioxygenase.

[0008] The positive pools were obtained from a cDNA library from chicken duodenum which was screened for β,β -carotene 15,15'-dioxygenase activity in a cellular transactivation assay. By this strategy several positive cDNA pools were identified. By combining the two strategies the gene coding for β,β -carotene 15,15'-dioxygenase could be successfully cloned. It is an object of the present invention to provide a protein having the vitamin A producing activity of β,β -carotene 15,15'-dioxygenase comprising an amino acid sequence which is identical or homologous to sequence ID no. 1 (shown in Fig. 4) whereby the degree of homology to sequence ID no. 1 is at least 60%.

[0009] After having provided the sequence of β,β -carotene 15,15'-dioxygenase isolated from chicken corresponding proteins from different animals like swine, cow, goat, dog, rabbit, poultry, fish and humans can easily be obtained. Since the sequence is known suitable regions of the nucleic acid sequence can be selected as primers for a polymerase chain reaction with a suitable nucleic acid which allows an easy and rapid amplification of the gene coding for the protein.

[0010] The present invention comprises therefore not only proteins having an amino acid sequence identical to the sequence given in sequence ID no. 1 but also such proteins which have an amino acid sequence homologous to the sequence ID no. 1. The degree of homology is, however, at least 60%, preferably 70%, more preferably 80% and especially preferred at least 90%. Homology as defined in the present invention means that when the amino acid sequences of two proteins are aligned at least the given percentage is identical. The alignment of the amino acids can be performed with the help of a suitable computer programme which is commercially available, preferably the Wisconsin Sequence Analysis Package GCG (Genetics Computer Group, University Research Park, Madison), Version 9.1, 1997. The remainder of the amino acids may be different. A homology of 90% for example means that 90% of the amino acids of the protein are identical compared with the amino acid sequence given in sequence ID no. 1 whereas 10% of the amino acids may be different. The proteins of the present invention have, however, the biological activity of β,β -carotene

[0011] 15,15'-dioxygenase which is explained above in more detail. Preferred proteins which are within the scope of the present invention are derived from other mammals or from humans.

[0012] Another aspect of the present invention concerns nucleic acid sequences coding for a protein having the biological activity of β,β -carotene 15,15'-dioxygenase. A nucleic acid sequence coding for the enzyme derived from chicken is shown in sequence ID no. 2 (see Figure 3). The nucleic acid sequences of the present invention code for a protein of the present invention or a part thereof. Shorter nucleotide sequences suitable for PCR have a length of at least 20 bases, preferably 25 bases and most preferred at least 30 bases.

[0013] The nucleic acid sequences of the present invention can be preferably used as primers for the specific amplification of a gene or part thereof coding for β,β -carotene 15,15'-dioxygenase. Primers can also be used for the specific amplification of 5' nontranslating or 3' nontranslating sequences of the cDNA described above. The nucleic acid sequences of β,β -carotene 15,15'-dioxygenase cDNA can be used as a probe for the detection of the coding as well as for the noncoding regions or parts thereof. The nucleic acid sequences of the present invention can be preferably used as antisense RNA probe for in situ hybridization.

[0014] It is especially preferred to use primers and probes having a part of the sequence given in sequence ID no. 2 as primers and/or probes in testkits which can be used for the amplification and/or detection of genes/mRNAs coding for β,β -carotene 15,15'-dioxygenase by polymerase chain reaction (PCR). The selection of suitable parts of the nucleic acid sequence can be performed by the person skilled in the art without difficulties. A nucleic acid sequence used as primer or probe is usually selected from a region which is highly conserved within the protein. Conserved means that the nucleic acid sequences of such regions of proteins obtained from different species are very similar. On the other hand the preferred nucleic acid sequence should not be present in other nucleic acid sequences which do not code for β,β -carotene 15,15'-dioxygenase, because this might lead to false positive results. By aligning several sequences derived from different species such regions can easily be determined. Although the nucleic acid can be a ribonucleic acid it is more preferred to have desoxyribonucleic acid sequences.

[0015] One preferred use in diagnostics is the determination of the β,β -carotene 15,15'-dioxygenase in patients. There is a variability in β -carotene cleavage potential among the population. Humans with low dioxygenase levels (with e.g. mutations or polymorphisms in the gene for β,β -carotene 15,15'-dioxygenase) could be identified and selected for vitamin A supplementation.

[0016] A diagnostic kit based on PCR can be designed to detect frequent mutations in the dioxygenase gene.

Another diagnostic option is quantification of mRNA by RT-PCR. With this diagnostic tool differences in expression of β,β -carotene 15,15'-dioxygenase in various tissues and in different species can be found.

[0017] Since the protein has been expressed and a method for purifying the protein is described in detail in the examples the person skilled in the art can use the protein or peptides derived from the amino acid sequences in order to generate antibodies which do specifically react with the protein. It is either possible to produce polyclonal antibodies by immunizing laboratory animals, like rabbits, sheeps or goats preferably with an adjuvant or monoclonal antibodies by the well-known technique described by Köhler and Milstein [European Journal of Immunology, 1976, 6 (7), p. 511-519]. The antibodies should specifically react with β,β -carotene 15,15'-dioxygenase in order to avoid an unspecific crossreaction. This means that the antibodies of the present invention should preferably react with an epitope which is present only on a protein of the present invention.

[0018] Such antibodies can be preferably used in immunoassays for the detection and/or quantification of β,β -carotene 15,15'-dioxygenase in a test fluid. The test fluid may be a liquid, like serum, obtained from a patient. There are several types of immunoassays which are well-known to the person skilled in the art. Very frequently one antibody, preferably a monoclonal antibody is fixed to a solid phase. This antibody is then brought into contact with the fluid containing the β,β -carotene 15,15'-dioxygenase and after washing it is further reacted with a second monoclonal antibody which binds to another epitope of the enzyme. The second antibody is usually labeled and shows the presence of the sandwich consisting of the antigen and two different antibodies.

[0019] The antibodies can also be used in laboratory methods like Western blots or immuno-precipitations. Preferably such antibodies can be used in immunohistochemistry to detect epitopes of β,β -carotene 15,15'-dioxygenase in embedded or fixed tissues or cells of any species of interest.

[0020] In a further embodiment of the present invention the β,β -carotene 15,15'-dioxygenase is used for the production of vitamin A whereby the enzyme cleaves enzymatically β -carotene into two molecules of retinal which will subsequently be reduced by retinol dehydrogenase to vitamin A. The β,β -carotene 15,15'-dioxygenase can be used to enzymatically convert β -carotene which may be obtained from plant sources. A preferred source of β -carotene is the alga *Dunaliella bardawil* which has a high endogenous level of β -carotene. Suitable algae can be grown conveniently and β -carotene can be purified therefrom at rather low cost. The carotene can be conveniently cleaved enzymatically by using a protein of the present invention. The carotene dioxygenase can preferably be immobilized in order to provide a continuous process.

[0021] Another aspect of the present invention concerns the introduction of the gene coding for a protein having β,β -carotene 15,15'-dioxygenase activity into a suitable host cell.

[0022] The first step in such a method is usually to insert the cDNA into a suitable vector. The vector must fit with the host cell into which the gene should be introduced. There are specific vectors available for bacteria, yeasts, plant cells, insect cells or mammalian cells. Preferably the gene is combined with genetic structures which provide the required genetic regulation like promoters, enhancers, ribosomal binding sites etc.

[0023] Systems for the expression of genes encoding carotenoid biosynthetic enzymes in procaryotes, especially in *E. coli* or *Bacillus subtilis* or *Flavobacter* and eucaryotes, e.g. fungi are known in the art and described e.g. in European Patent Application Publication No. (EP) 747 483 or in EP 872 554.

[0024] The vector having the gene and the other required genetic structures is then introduced into suitable host cells by well-known methods like transformation, transfection, electroporation or microprojectile bombardment. Depending on the host cell it may be preferred to stably integrate the gene coding for a protein of the present invention into the genome of the host cell. The cells obtained by such methods can then be further propagated and if the cell is a plant cell it is possible to generate therefrom transgenic plants. In one embodiment of the present invention host cells are plant cells and tomato cells are especially preferred. The technology to produce transgenic tomatoes is well-established and the tomato contains sufficient β -carotene in order to come up with a reasonable vitamin A level after introduction of carotene dioxygenase into the tomato plant. In green pepper, melon or especially carrot the endogenous level of β -carotene is even higher and therefore also these plants are especially preferred.

[0025] Another preferred embodiment of the present invention concerns algae. Halotolerant algae may contain high levels of β -carotene. A transfection of such algae with an expression vector comprising the β,β -carotene 15,15'-dioxygenase cDNA leads to a high intracellular vitamin A level which can easily be recovered from such algae by simple purification steps.

[0026] In another aspect of the present invention a gene coding for a protein of the present invention can be introduced into mammalian cells and especially into human cells. It is for example possible to insert the gene coding for a β,β -carotene 15,15'-dioxygenase into suitable cells, for example peripheral blood stem cells. Such cells which contain the gene for β,β -carotene 15,15'-dioxygenase may be administered to people having mutations or deletions in the β,β -carotene 15,15'-dioxygenase gene. Such mutations and deletions, respectively, may have the effect that such patients are not able to cleave β -carotene enzymatically. Therefore, such patients always have a low vitamin A level and thus suffer from various developmental and ophthalmological problems. The administration of suitably transfected cells expressing the β,β -carotene 15,15'-dioxygenase to such patients by way of somatic gene therapy is a promising way to

improve their situation.

[0027] Some of the most important results of the present invention are summarized in the Figures.

[0028] Figure 1 shows the result from the last step of purification of β,β -carotene 15,15'-dioxygenase from the small intestine of chicken. The SDS-PAGE pattern and β,β -carotene 15,15'-dioxygenase activity of individual fractions from the gel permeation chromatography run are shown. On the gel, the protein A is marked by an arrow and seemed to correlate best with the β,β -carotene 15,15'-dioxygenase activity. It was therefore chosen for further amino acid sequence analysis. The abbreviations have the following meaning: Std.: molecular weight standard; conc.: concentrate loaded onto the gel permeation chromatography column.

[0029] Figure 2 shows schematically the transactivation assay in eukaryotic cells. cDNAs are transfected and expressed in MCF-7 cells. When incubated with β -carotene, a positive pool shows cleavage activity. The cleavage product retinal is further oxydized to retinoic acid (RA) which binds to the endogenous receptor. The receptor/ligand complex binds to the response element on the reporter plasmid and leads to an enhanced transcription of the luciferase gene. The luminescence signals are detected in a luciferase assay with a sensitive CCD camera.

[0030] Figure 3 shows the cDNA sequence (sequence ID no. 2) for β,β -carotene 15,15'-dioxygenase which has a length of 3090 base pairs excluding the poly A tail. 132 base pairs are 5' nontranslating sequence, the coding sequence has 1578 base pairs and the 3' nontranslating sequence 1380 base pairs, respectively. A putative poly A signal is found at position 3073.

[0031] Figure 4 shows the derived amino acid sequence (sequence ID No. 1) of β,β -carotene 15,15'-dioxygenase derived from chicken having 526 residues. The amino acid sequence is given in the one letter code.

[0032] Figure 5 shows a comparison of the β,β -carotene 15,15'-dioxygenase amino acid sequence with a protein having the designation RPE65 which was found by a sequence comparison in EMBL Genbank as the protein having the highest homology to the β,β -carotene 15,15'-dioxygenase of the present invention.

[0033] Figure 6 shows two fractions of β,β -carotene 15,15'-dioxygenase eluted from a Co^{2+} -chelate column. In lanes 1 and 2 two different fractions were loaded and lane 3 is a low range molecular weight marker.

[0034] Figure 7 shows an HPLC analysis of an activity test of β,β -carotene 15,15'-dioxygenase which was cloned and expressed in *E. coli*.

[0035] Figure 8 is a proof that the peak representing the only product of the enzymatic cleaving is retinal.

[0036] The present invention is further illustrated in the following examples:

Example 1:

Assay of β,β -carotene 15,15'-dioxygenase activity

[0037] For the tests the following solutions were prepared:

- a) Solution 1 (mixed micelle solution): Glycocholic acid (1.16 g) was dissolved in 5 ml H_2O under stirring and by drop-wise addition of 5 N NaOH. After pH was adjusted to 6.8-7.2 with acetic acid and the volume increased to 10 ml with H_2O , 80 mg of asolectin (Fluka) were added and dissolved under stirring.
- b) Solution 2 (substrate solution): 500 μl of an α -tocopherol solution (10 mg/ml in hexane) and 235 μg of a β -carotene solution (80 $\mu\text{g}/\text{ml}$ of pure all-E- β -carotene in benzene) were mixed in a glass vial, protected from light and the solvents evaporated under a gentle nitrogen stream. 1 ml of solution 1 was added under vortexing and eventually a few short ultrasonic bursts until a clear solution occurred.
- c) Solution 3 (homogenization buffer): 100 mM KH_2PO_4 adjusted with 5 N KOH to pH 7.8 and containing 4 mM MgCl_2 , 6 H_2O and 30 mM nicotinamide.
- d) Solution 4 (GSH solution): 60 mg/ml reduced glutathione dissolved in solution 3.
- e) Solution 5 (standard solution): 10 $\mu\text{g}/\text{ml}$ vitamin A acetate in hexane/chloroform 9:1.

[0038] Activity assay: 2 ml enzyme preparation (approx. 4 mg protein, assayed by BCA protein assay, Pierce Chemicals) were placed in a light-protected glass vial in a shaking water bath (30 min., 37 °C). 0.2 ml solution 4 was added and the reaction was started after 2 min. of temperature equilibration by addition of 50 μl solution 2. After 3 h, the reaction was stopped by placing the vials on ice and subsequent addition of 1 ml acetonitrile followed by 5 ml chloroform. The vials were vortexed 3 times for 7 s and phase separation was enforced by centrifugation for 5 min. at 5000 g. Extraction was repeated twice with 0.6 ml chloroform. The combined chloroform phases were evaporated and resolubilized in 200 μl solution 5 under short sonication. Insoluble material was removed by filtration through 0.45 μm filters. An aliquot of 20 μl was separated by HPLC on a reversed phase C_{18} column (Lichrospher 100, 5 μm , 12.5 cm x 4.6 mm, Bischoff Chromatography, Leonberg, Germany; 1 ml/min, column temperature 25°C) with a discontinuous, optimized gradient of acetonitrile/tetrahydrofuran/(1% ammonium acetate in H_2O) from 50:20:30 (eluent A) to 50:44:6 (eluent B). These conditions allow complete separation of β,β -carotene and retinal as well as apo- β -carotenals and retinoic

acids. Calibration curves were made for both β,β -carotene and retinal in the concentration ranges 2-40 and 1-10 ng/ μ l, respectively, and were correlated to the value of vitamin A acetate which served as an internal standard. Enzymatic activity was expressed as the amount of retinal liberated in the activity assay during 3 h of incubation at 37 °C (100% = 17.6 nmol).

Example 2:

Purification of β,β -carotene 15,15'-dioxygenase

[0039] Purification was done as rapidly as possible, and all buffers and equipments were cooled to 4 °C.

[0040] Solution 6 (protease inhibitor-containing homogenisation buffer): 125 mM benzamidine \cdot HCl, 250 mM 6-aminocaproic acid and 125 μ M soybean trypsin inhibitor were dissolved in H₂O by sonication. A 4 ml-aliquot of this solution was mixed with 100 ml of solution 3.

[0041] Solution 7: 10 mM KH₂PO₄, 1 mM reduced glutathione, pH 7.8.

[0042] Solution 8 (eluent A, phenyl-Sepharose chromatography): 10 mM KH₂PO₄, 1 mM reduced glutathione, 0.5 M (NH₄)₂SO₄, pH 7.8.

[0043] Solution 9 (eluent B, phenyl-Sepharose chromatography): 10 mM KH₂PO₄, 1 mM reduced glutathione, 10% glycerol, pH 7.8.

[0044] Solution 10 (eluent B, Poros HQ chromatography): 10 mM KH₂PO₄, 1 mM reduced glutathione, 0.5 M NaCl, 10% glycerol, pH 7.8.

[0045] Solution 11 (elution buffer for gel permeation chromatography): 50 mM KH₂PO₄, 1 mM reduced glutathione, 150 mM NaCl, 10% glycerol, pH 7.8.

[0046] Laying hens at age of 20-24 weeks (strain Lohmann LSL, Hatchery Wuethrich, CH-3123 Belp, Switzerland) were kept on a pigment-free chicken diet (Kliba 3179, Kliba, CH-4303 Kaiseraugst, Switzerland). The animals were killed by decapitation and the first 20 cm of the duodenal loop was removed, separated from pancreas and rinsed with 40 mL each of 0.9% NaCl solution. The intestines were immediately frozen in dry ice and stored at -80°C until use.

[0047] Ten intestines were thawed on ice in approx. 2 h and opened length-wise in an ice cooled Petri dish. The mucosa was scraped off with a slide, weighed and homogenized in a Teflon-glass Potter-Elvehjem homogeniser in 4 volumes of solution 6 with six strokes. Upon centrifugation at 62000 g for 1 h, the clear supernatant was divided into 32 aliquots of 15 ml each. From these preparations an ammonium sulphate fractionation was made. The precipitate obtained from the 20-45% step was centrifugated at 5000 g for 10 min. and the pellet was stored at -80°C for further use.

[0048] Ten aliquots of the (NH₄)₂SO₄ pellet were dissolved in 150 ml of solution 7, sterile-filtered and loaded on a HiLoad 26/10 phenyl-Sepharose High Performance column (column volume 53 ml; Pharmacia, Uppsala, Sweden) equilibrated with solution 8. Proteins were eluted at a flow rate of 8 ml/min with a steep gradient over 1 column volume (CV) from solution 8 to solution 9. β,β -Carotene 15,15'-dioxygenase eluted at a conductivity of < 15 mS/cm, but only fractions with a conductivity of < 1 mS/cm were pooled and directly loaded onto a 30 ml Blue Sepharose 6 Fast Flow column (Pharmacia) equilibrated with solution 9. β,β -Carotene 15,15'-dioxygenase activity eluted (at a flow rate of 8 ml/min) in the break-through fractions which were (again) directly loaded onto a 20 ml Poros HQ/M anion exchange chromatography column (PerSeptive Biosystems, Framingham, MA, USA) equilibrated with solution 9. β,β -Carotene 15,15'-dioxygenase was eluted at a flow rate of 15 ml/min with a linear gradient over 18 CV from solution 9 to solution 10. Activity was detected in the gradient in a conductivity range of 10-20 mS/cm. The pooled fractions (70 ml) were concentrated to ~1.3 ml with Ultrafree-15 filter units (MW cut-off 50,000; Millipore, Bedford, MA, USA). An aliquot of the concentrate (500 μ l) was loaded onto a Superdex 200 HR 10/30 gel filtration column (CV 24 ml; Pharmacia) and eluted at a flow rate of 0.5 ml/min with solution 11. Aliquots of each fraction were used for activity assays (see example 1) and, upon concentration, for SDS-PAGE (with MOPS running buffer) on 10% NuPAGE gels (Novex, San Diego, CA, USA). The results of this experiment are shown in Fig. 1 and Table 1.

Purification step	Total protein [mg]	Total activity [nmol/h]	Yield [%]	Specific activity [pmol/(h·mg)]	Purification factor
(NH ₄) ₂ SO ₄ pellet	779	8.61	100	11.0	—
phenyl-Sepharose	80.8	8.27	96.1	102	9.27
Blue	16.0	8.86	103	554	50.1
Sepharose Poros HQ	1.56	3.90	45.3	2500	226

Table 1: Summary table for the purification of β,β -carotene 15,15'-dioxygenase starting with 10 aliquots of the (NH₄)₂SO₄ pellet (means of 3-4 measurements).

Example 3:

Amino acid sequence information for β,β -carotene 15,15'-dioxygenase

[0049] For amino acid sequence analysis, fractions of the gel filtration run (as shown in Fig. 1) were separated on a 8-16% Tris/glycine gel (Novex), and the proteins transferred to an Immobilon P^{SO} membrane (Millipore) and stained with amido black.

[0050] Since protein A proved to be N-terminally blocked, multiple aliquots of fraction 18 from the gel filtration run (see Fig. 1) were separated on a 10% Tris/glycine gel (Novex), and the gel was stained with Colloidal Coomassie Blue (Novex). The band corresponding to protein A was excised from the gel, and the protein digested in-gel with trypsin. The tryptic digest was separated by micro-bore reversed-phase HPLC on a 150 x 1.0 mm Vydac C₁₈ column (Vydac, Hesperia, CA, USA). Peptides were eluted with an acetonitrile gradient in 0.1% trifluoroacetic acid, and peptide containing fractions were collected for further analysis. Two fractions could be identified by MALDI-TOF-MS (Voyager Elite, PerSeptive Biosystems) to contain one single peptide each. N-terminal Edman degradation revealed the following sequences:

(1) Ala-Glu-Val-Gln-Gly-Gln-Leu-Pro (Seq. ID No.3)

(2) Asn-Lys-Glu-Glu-His-Pro-Glu-Pro-Ile-Lys-Ala-Glu-Val-Gln-Gly-Gln-Leu-Pro (Seq. ID No. 6)

[0051] Note that the last 8 amino acids of peptide (2) correspond to peptide (1).

Example 4:

Cloning the full length cDNA for the β,β -carotene 15,15'-dioxygenase

A) RNA isolation:

[0052] 4 week old Vedette chicken were killed, the duodenum was removed, washed with sterile PBS and cut open with scissors. The mucosal layer was scraped off with a glass slide, weighed and homogenized immediately with a Polytron in 1 ml of Trizole reagent (Life Technologies) per 100 mg of tissue. Then the standard protocol from Life Technologies was followed. Poly A mRNA was isolated by the polyATtract mRNA Isolation kit from Promega Corporation, Madison.

B) PCR and RT-PCR:

[0053] In the peptide sequence NKEEHPEPIKAEVQGQLP (peptide 2 of Example 3) (Seq. ID No. 7) two degenerate primers were designed: In order to have a lower degeneracy the base Inosin was used in one and in two wobble positions, respectively.

5' primer: 5'AAC AAR GAR GAS CAY CCI GA 3' (Seq. ID No. 8) (20 mer with a degeneracy of 16x)

3' primer: 5'SAG CTG ICC CTG IAC YTC SGC 3' (Seq. ID No. 9) (21 mer with a degeneracy of 8x)

R= A or G, S= C or G, Y= C or T

[0054] The oligos were synthesized on a Pharmacia Gene Assembler Plus using standard phosphoramidite chemistry. Deprotection was done with 1 ml conc. ammonium hydroxide solution (Applied Biosystems) and final desalting was performed with a NAP 10 column (Amersham Pharmacia Biotech).

[0055] For PCR 100 ng chicken duodenal cDNA were taken as template and the following steps performed: 94°C 30"; 52°C 30"; 72°C 1' for 40 cycles. The resulting band of 51 bp was cut out from a 10% polyacrylamide gel, electrophoresed on DEAE paper at 300 V for 1.5 h, eluted from the DEAE paper once with 40 µl and twice with 30 µl 1.5 M NaCl, 5 mM Tris, 0.5 mM EDTA, precipitated with 2.5 volumes of ethanol 100% and 1 µg glycogen, washed with 0.5 ml of 80% ethanol, dried and dissolved in 20 µl TE (10 mM Tris, 1 mM EDTA).

[0056] The resulting fragment of 51 bp was cloned into pGEM-T Easy, a commercially available T/A cloning vector (Promega Corporation, Madison). The corresponding cDNA sequence was determined by automated fluorescent sequencing on a Vistra DNA Sequencer 725 (Amersham Pharmacia Biotech).

[0057] From the above DNA sequence a homologous forward primer was derived:

5' TCTGAATTCGGAGCCCATAAAAGC 3' (primer dioxy12) [Seq. ID No. 10]

[0058] At the 5' end an EcoRI site (underlined sequence) was introduced; the following 17 nucleotides are homologous to the previously obtained dioxygenase sequence.

[0059] In a RT-PCR reaction a polyT/Not primer (commercially available from Invitrogen, San Diego) was used as reverse primer together with primer dioxy12.

[0060] One tube RT-PCR kit from Boehringer Mannheim was taken and the corresponding protocol followed:

mix 1:	
18.3 µl	H ₂ O
2.5 µl	DTT (100 mM)
1.0 µl	dNTPs (10 mM)
1.0 µl	oligo dT/Not (0.2 µg/µl) (3' primer)
1.0 µl	dioxy12 (5' primer) (20 µM)
0.2 µl	RNAse inhibitor (40 U/µl)
1.0 µl	chicken duodenal total RNA (2.2 µg/µl)
25.0 µl	
mix 2:	
14.0 µl	H ₂ O
10.0 µl	RT-PCR buffer 5x
1.0 µl	enzyme mix (AMV RT, Taq and Pwo DNA Polymerase)
25.0 µl	

[0061] The 2 mixes were combined and the PCR protocol started on a MJ Research PTC200 DNA Engine.

50°C	30'	
94°C	2'	
94°C	30"	
57°C	30"	10 cycles
68°C	45"	
94°C	30"	
62°C	30"	25 cycles
68°C	45" + 3"	/cycle
68°C	7'	
4°C	over night	

[0062] With this RT-PCR protocol a band of 597 bp was amplified from chicken total duodenal RNA. The PCR band was isolated from a 1% agarose gel, cloned into pGEM-T Easy cloning vector and subsequently sequenced. The original peptide is present in the sequence as well as an open reading frame over the whole sequence of 597 bp.

C) Chicken cDNA-library:

[0063] From chicken duodenal polyA⁺ RNA cDNA was made with the Copy Kit (Invitrogen, San Diego) using a modified Gubler-Hoffman procedure. The cDNA was size-selected (0.9-5.5 kb) and cloned into the eukaryotic expression vector pcDNA1.1/Amp (Invitrogen).

[0064] Electroporation into E.coli Top10 was done with a Bio-Rad Gene Pulser II system following the standard protocol. This resulted in a cDNA library of 480'000 individual clones. The library was splitted into 250 pools with 1500-2500 individual clones each. Each pool was amplified in 100 ml LB medium: Bacterial growth was stopped at OD 0.8-1.0 by adding chloramphenicol to a final concentration of 170 µg/ml. Incubation was continued over night in order to increase the DNA amount.

D) Activity screening of the chicken cDNA library:

[0065] 90 of the above pools were tested for activity in a transactivation assay based on the detection of retinoic acid which is produced in eukaryotic cells after β-carotene cleavage. The principle of the activity test is shown in Fig. 2.

[0066] 5 µg of DNA from each pool were transfected with 20 µg of lipofectin (Life Technologies) into a reporter cell line bearing a luciferase reporter plasmid with a RARE (retinoic acid response element) in front of the tk promoter (Herpes simplex thymidine kinase promoter). Transfections were done for 7 h under serum free conditions. After 7 h the transfection mix was removed and RPMI medium with 10% charcoal treated FCS (fetal calf serum) was added. After 20 h of incubation β-carotene (β-carotene 10% CWS, F. Hoffmann-La Roche Ltd.) or a placebo formulation were added to the culture medium to a final β-Carotene concentration of 5 µM. Incubation was continued for 18 h. Then cells were washed with PBS, and luciferase expression was measured after substrate addition with a nitrogen cooled slow scan CCD camera (AstroCam Ltd.) Exposure time usually was 8 min. Analysis was done with the Image Pro Plus 3.0 software package (Media Cybernatic, Maryland). 3 pools were strongly positive, 7 pools showed weaker, but detectable activity.

[0067] One of the positive pools was plated on a square agar plate. 2 filters (nylon membranes, Gene Screen, NEN Research Products, Boston) were processed and screened with the radioactively ([α³²P]dATP, Amersham) labeled 597 bp PCR-fragment. From 9500 colonies screened, 14 were double positive. From 36 colonies picked, 5 showed the same pattern after restriction site analysis. 2 clones were sequenced from the 5' end and the original 51 bp sequence was found. Subsequently the whole cDNA was sequenced and confirmed twice.

[0068] All molecular biological procedures were done according to Sambrook, Fritsch and Maniatis, Laboratory Manual, Cold Spring Harbor Laboratory Press, 1989, if not otherwise mentioned.

[0069] The obtained cDNA sequence is shown in Figure 3 and the amino acid sequence deduced therefrom in Figure 4.

[0070] Figure 4 shows the derived amino acid sequence having 526 residues.

Example 5: Sequence comparison:

[0071] By sequence comparison with the EMBL Genbank a high homology between the known protein RPE65 (Hamel et al., J.Biol.Chem. [1993] p. 15751-15757) and the β , β -carotene 15,15' dioxygenase was found. A homology of 55.5% on the aminoacid level was found. The sequence alignment is shown in Figure 5.

Example 6:

Expression of the cDNA for β , β -carotene 15,15'-dioxygenase in E. coli:

[0072] With PCR the coding sequence of the β , β -carotene 15,15'-dioxygenase cDNA was amplified and the resulting fragment of 1578 bp was cloned into the EcoRI/BamHI site of the prokaryotic expression vector pQE-12 (Qiagen). The vector contains an in frame hexa-His affinity tag at the C-terminus of the dioxygenase.

[0073] In addition this plasmid contains a regulated promoter with two lacI repressor binding sites.

[0074] The E.coli strain M15pREP4 was transformed with the expression plasmid. For expression 1 l LB medium, containing 100 μ g/ml ampicillin and 25 μ g/ml kanamycin, was inoculated with 30 ml of an overnight culture. Growth was allowed until OD₆₀₀ of 0.6-0.8. At this point the culture was induced with 1 mM IPTG (isopropyl- β -thiogalactoside) and growth continued for another 1.5-2 hours. Bacteria were harvested by centrifugation and the pellet was frozen at -80°C.

[0075] The pellet was thawed by stirring in 20 ml extraction buffer (300 mM NaCl, 50 mM NaH₂PO₄, 20 mM Tris-HCl, 1 mg/ml Tween 40; pH 7.8) including 2.5 mg/ml dodecyl- β -D-maltoside. 1 ml protease inhibitor cocktail (5.9 mM benzamidine-HCl, 10 mM 6-amino-caproic acid, 5 μ M soybean trypsin inhibitor) was added at the same time.

[0076] The bacterial cells were lysed by 4 min. treatment with a Polytron (Kinematica AG, Switzerland) using a PT7 unit following 4 min. sonication with a Branson Sonifier 250.

[0077] The lysate was spun at 12000 x g and the supernatant purified over a Co²⁺-chelate column (Talon Superflow Resin, Clontech, Heidelberg, Germany). The protein was eluted with 15-20 ml of extraction buffer containing 150 mM imidazole.

[0078] The fractions were loaded on a 10% polyacrylamide gel and the protein containing fractions dialysed against 150 mM tricine, 5 mM FeSO₄, 3 mg/ml reduced glutathione, 0.21 mg/ml sodium cholate. The gel is shown in Fig. 6. The samples were then assayed in the dioxygenase activity assay.

Example 7:

Expression of recombinant β , β -carotene 15,15'-dioxygenase in the human duodenal cell line HuTu80:

[0079] With PCR the coding sequence of the β , β -carotene 15,15'-dioxygenase cDNA was amplified and the resulting fragment of 1578 bp was cloned into the BamHI/XhoI site of the plasmid pSFV₂gen. This vector is part of the Semliki Forest Virus expression system, which works highly efficiently in most mammalian cells.

[0080] The plasmid was used for in vitro synthesis of recombinant RNA, which was subsequently electroporated together with a helper virus into BHK cells (baby hamster kidney cells) for production of a high titer virus stock. With an aliquot of this stock the human duodenal cell line HuTu80 was infected. 16-18 h after infection the cells were harvested and the pellet frozen at -80°C. Either the whole cell pellet, the cytoplasmic fraction or the membrane fraction was used in an activity assay. β , β -carotene 15,15'-dioxygenase activity was found in the whole cell extract and in cytosolic fractions, while in the membrane fraction no activity was detected.

Example 8:

β , β -carotene 15,15'-dioxygenase activity of the recombinant protein expressed in E.coli and in human cells:

[0081] After expression in E. coli and purification over a metal chelate column, the protein shows cleavage activity with β -carotene as substrate. Retinal was the only product detected by HPLC after incubation with β -carotene. No apocarotenals or other metabolites were found. This could be proved by HPLC analysis as shown in Fig. 7 and 8.

SEQUENCE LISTING

5

<110> F. Hoffmann La Roche AG

10

<120> beta,beta-carotene 15,15'-dioxygenases, nucleic acid
sequences coding therefore and their use

<130>

<140>

15

<141>

<150> 99.103382.0

<151> EP 1999-02-22

20

<160> 10

<170> PADAT Sequenzmodul, Version 1.0

25

30

35

40

45

50

55

EP 1 031 627 A1

<210> 1
 <211> 526
 <212> PRT
 <213> Gallus gallus

<400> 1

Met Glu Thr Ile Phe Asn Arg Asn Lys Glu Glu His Pro Glu Pro Ile
 1 5 10 15
 Lys Ala Glu Val Gln Gly Gln Leu Pro Thr Trp Leu Gln Gly Val Leu
 20 25 30
 Leu Arg Asn Gly Pro Gly Met His Thr Ile Gly Asp Thr Lys Tyr Asn
 35 40 45
 His Trp Phe Asp Gly Leu Ala Leu Leu His Ser Phe Thr Phe Lys Asn
 50 55 60
 Gly Glu Val Tyr Tyr Arg Ser Lys Tyr Leu Arg Ser Asp Thr Tyr Asn
 65 70 75 80
 Cys Asn Ile Glu Ala Asn Arg Ile Val Val Ser Glu Phe Gly Thr Met
 85 90 95
 Ala Tyr Pro Asp Pro Cys Lys Asn Ile Phe Ala Lys Ala Phe Ser Tyr
 100 105 110
 Leu Ser His Thr Ile Pro Glu Phe Thr Asp Asn Cys Leu Ile Asn Ile
 115 120 125
 Met Lys Thr Gly Asp Asp Tyr Tyr Ala Thr Ser Glu Thr Asn Phe Ile
 130 135 140
 Arg Lys Ile Asp Pro Gln Thr Leu Glu Thr Leu Asp Lys Val Asp Tyr
 145 150 155 160
 Ser Lys Tyr Val Ala Val Asn Leu Ala Thr Ser His Pro His Tyr Asp
 165 170 175
 Ser Ala Gly Asn Ile Leu Asn Met Gly Thr Ser Ile Val Asp Lys Gly
 180 185 190
 Arg Thr Lys Tyr Val Leu Phe Lys Ile Pro Ser Ser Val Pro Glu Lys
 195 200 205
 Glu Lys Lys Lys Ser Cys Phe Lys His Leu Glu Val Val Cys Ser Ile
 210 215 220
 Pro Ser Arg Ser Leu Leu Gln Pro Ser Tyr Tyr His Ser Phe Gly Ile
 225 230 235 240
 Thr Glu Asn Tyr Ile Val Phe Ile Glu Gln Pro Phe Lys Leu Asp Ile
 245 250 255
 Val Lys Leu Ala Thr Ala Tyr Ile Arg Gly Val Asn Trp Ala Ser Cys
 260 265 270
 Leu Ser Phe His Lys Glu Asp Lys Thr Trp Phe His Phe Val Asp Arg
 275 280 285
 Lys Thr Lys Lys Glu Val Ser Thr Lys Phe Tyr Thr Asp Ala Leu Val
 290 295 300
 Leu Tyr His His Ile Asn Ala Tyr Glu Glu Asp Gly His Val Val Phe
 305 310 315 320
 Asp Ile Val Ala Tyr Arg Asp Asn Ser Leu Tyr Asp Met Phe Tyr Leu
 325 330 335
 Lys Lys Leu Asp Lys Asp Phe Glu Val Asn Asn Lys Leu Thr Ser Ile
 340 345 350
 Pro Thr Cys Lys Arg Phe Val Val Pro Leu Gln Tyr Asp Lys Asp Ala
 355 360 365

EP 1 031 627 A1

	Glu	Val	Gly	Ser	Asn	Leu	Val	Lys	Leu	Pro	Thr	Ser	Ala	Thr	Ala	Val
	370						375					380				
5	Lys	Glu	Lys	Asp	Gly	Ser	Ile	Tyr	Cys	Gln	Pro	Glu	Ile	Leu	Cys	Glu
	385					390					395					400
	Gly	Ile	Glu	Leu	Pro	Arg	Val	Asn	Tyr	Asp	Tyr	Asn	Gly	Lys	Lys	Tyr
					405					410					415	
	Lys	Tyr	Val	Tyr	Ala	Thr	Glu	Val	Gln	Trp	Ser	Pro	Val	Pro	Thr	Lys
				420					425					430		
10	Ile	Ala	Lys	Leu	Asn	Val	Gln	Thr	Lys	Glu	Val	Leu	His	Trp	Gly	Glu
		435					440						445			
	Asp	His	Cys	Trp	Pro	Ser	Glu	Pro	Ile	Phe	Val	Pro	Ser	Pro	Asp	Ala
	450						455					460				
	Arg	Glu	Glu	Asp	Glu	Gly	Val	Val	Leu	Thr	Cys	Val	Val	Val	Ser	Glu
15	465					470					475					480
	Pro	Asn	Lys	Ala	Pro	Phe	Leu	Leu	Ile	Leu	Asp	Ala	Lys	Thr	Phe	Lys
					485					490					495	
	Glu	Leu	Gly	Arg	Ala	Thr	Val	Asn	Val	Glu	Met	His	Leu	Asp	Leu	His
				500					505					510		
20	Gly	Met	Phe	Ile	Pro	Gln	Asn	Asp	Leu	Gly	Ala	Glu	Thr	Glu		
		515						520					525			

25

30

35

40

45

50

55

EP 1 031 627 A1

<210> 2
 <211> 3111
 <212> DNA
 <213> Gallus gallus

<220>
 <223>

<400> 2

cggtaccact agtaacggcc gccagtgtgg tggaatccat ccttctatgt aacaggaaag 60
 agctgttctt agcccagaga ggagggcacc gtacgcctgc aggagcagct gggtagagga 120
 cacaggagag cgatggagac aatatttaac agaaacaaag aagagcatcc agagcccata 180
 aaagctgagg tgcaaggcca gttgccctact tggttgcaag gggtaacttct ccgaaatggc 240
 ccaggggatgc acacaatagg ggacactaaa tacaaccact ggtttgatgg cttggctctg 300
 ctgcacagct tcacgtttaa aaatggtgaa gtttactaca gaagtaagta cctccgaagt 360
 gacacataca actgcaatat agaagcaaac cgaatcgtgg tgtctgagtt tggaaccatg 420
 gcttatccgg atccatgcaa aaacatattt gccaaaggcat tctcatactt atctcacacc 480
 attcctgagt tcacggacaa ctgcctgac aacattatga aaactgggga tgattattat 540
 gctaccagtg agactaactt catcagaaaa attgatccac agactctgga gacactagat 600
 aaggtagact acagcaata thtagctgta aacttggcaa cttctcacc acactatgac 660
 agtgctggaa atattctcaa catgggtact tcaattgttg ataaaggag aacaaaatat 720
 gttctcttta agatcccttc ctctgtacca gaaaaagaaa agaagaaatc ttgttttaaa 780
 cacctggaag tagtatgctc catcccttct cgctccctgc tccaaccaag ctactaccac 840
 agctttggaa tcacagaaaa ttatatgttg ttcataagac agccatttaa actggatatt 900
 gtcaaaactgg caactgccta catccgaggt gtgaactggg cttcctgcct ttcctttcat 960
 aaggaggata agacgtggtt tcactttgta gacagaaaga cgaaaaaaga agtatccacc 1020
 aagttttaca ctgatgcttt ggtgctttat caccacataa atgcttacga agaagatggc 1080
 cacgttggtt ttgatatcgt tgcctacaga gacaatagct tgtacgatat gttttactta 1140
 aaaaaactgg acaaagactt tgaagtgaac aacaagctta cctccatccc aacctgcaag 1200
 cgctttgttg tgcctctgca gtatgacaag gatgcagaag taggttctaa tttagtcaaa 1260
 cttccaactt ccgcaactgc tgtaaaaga aaagatggca gcctctattg tcaacctgaa 1320

EP 1 031 627 A1

atattatgtg aagggataga actgectcgt gtcaactatg actacaatgg caaaaaatac 1380
 5 aagtatgtct atgcaacaga agtccagtgg agcccagttc ctacaaagat tgcaaaaactg 1440
 aatgtccaaa caaaggaagt actgcactgg ggagaagacc actgctggcc ctcagagccc 1500
 atctttgttc ccagccccga tgcaagagaa gaggatgaag gtgttgtttt gacctgtgtt 1560
 10 gtggtgtctg agccaaataa agcacccttc ctactcatct tggatgctaa aacattcaaa 1620
 gaattggggc gagccacagt taacgtagaa atgcatctgg acctgcatgg gatgtttata 1680
 ccacagaatg atttgggggc tgagacggaa taaaacgcta ttgatccgac tacacaaact 1740
 15 gagacaactt tctactgaac atgagttaat atccctttta ccattcaaga acaaccatat 1800
 aacgacacaa aatgactatg tataatctct taaataatag atataatcct ttaaggcac 1860
 agcgatgagt tttactacag gtaacgatat gcacaactgg catataacta ttccaaaaga 1920
 20 agaagaacga tcagtgtttt agaagtgtca atgttgtaga taacggcggc agaggggaaca 1980
 ggagagaaag gtaacgggaa tatttaatag aatatagatt tctgagcaaa tgaagtgcag 2040
 25 tatttatggg gtgatgcag gcgatgagca cataggtctg cagctcatgt atctttttaga 2100
 gatcgtttca agattgcagc ttgtgatgca agttttctcc agccagaaaa cctcatttta 2160
 aaccatctgc tactggtaat tcataccaat gcattttctt ggtgctcgat ttacactata 2220
 30 accaaagtta agtattacat tcaggtgcta caactttcta atttacaacc gaaacaaaca 2280
 agcaaacagc acttgctttg ctaataaccc catggtgtat ttttcctttt tatgatgaca 2340
 aaaccaagta catatggttt tatgtagcat tcaattatac ttcagtgcta ttccatccta 2400
 35 atgttataag caatttgtat ttaantcagt ttctcttgag aatatctgac ataacatttt 2460
 gtgtaatgag atgactatgt tgtctaaaga tgaacaggaa tgtatctttt attagtattg 2520
 ttaattgtgt tactaatact atgcatatga atgagagcaa tgtatttcta ggagaactca 2580
 40 gatatacatt caacaatttc tgtaggtgaa aatgcattta ctgatgaaag ttgaatcggt 2640
 aatgagggag aaaactgggt atccatccat ccaactatgt taggtgttca cctggtctgt 2700
 45 atgtgacacc acgctgtttg ggtatctctc actttccat acctgttctc atggtttctg 2760
 ctactcactg tattttgcag gagaganaca aaatgaaac actgtcactt actatcgccc 2820
 catcacataa gaacaatggg gctttggtga cttgttcatg attacataag atgtttgcag 2880
 50 cagagcagca atagaaccaa caccatccac agttcttgct tgctctgtta tgactccctt 2940
 tgctgtcttt atggtttgca tgtatgaaga atacactgcc taattctaant gtaaaaaagt 3000

55

EP 1 031 627 A1

cactggggtc agatctagag ctttaagtaag cagtctgggg ttttcaaagt tttatatggt 3060

ccataaaatg gaaataaaca cctccataat aaaaaaaaaa aaaaaaaaaa a 3111

<210> 3
<211> 8
<212> PRT
<213> Gallus gallus
<400> 3

Ala Glu Val Gln Gly Gln Leu Pro
1 5

<210> 4
 <211> 506
 <212> PRT
 <213> Gallus gallus

<400> 4

10	Glu	Glu	His	Pro	Glu	Pro	Ile	Lys	Ala	Glu	Val	Gln	Gly	Gln	Leu	Pro
	1				5					10					15	
	Thr	Trp	Leu	Gln	Gly	Val	Leu	Leu	Arg	Asn	Gly	Pro	Gly	Met	His	Thr
			20						25					30		
	Ile	Gly	Asp	Thr	Lys	Tyr	Asn	His	Trp	Phe	Asp	Gly	Leu	Ala	Leu	Leu
		35					40						45			
15	His	Ser	Phe	Thr	Phe	Lys	Asn	Gly	Glu	Val	Tyr	Tyr	Arg	Ser	Lys	Tyr
	50						55					60				
	Leu	Arg	Ser	Asp	Thr	Tyr	Asn	Cys	Asn	Ile	Glu	Ala	Asn	Arg	Ile	Val
	65				70						75					80
	Val	Ser	Glu	Phe	Gly	Thr	Met	Ala	Tyr	Pro	Asp	Pro	Cys	Lys	Asn	Ile
20					85					90					95	
	Phe	Ala	Lys	Ala	Phe	Ser	Tyr	Leu	Ser	His	Thr	Ile	Pro	Glu	Phe	Thr
				100					105					110		
	Asp	Asn	Cys	Leu	Ile	Asn	Ile	Met	Lys	Thr	Gly	Asp	Asp	Tyr	Tyr	Ala
		115					120					125				
25	Thr	Ser	Glu	Thr	Asn	Phe	Ile	Arg	Lys	Ile	Asp	Pro	Gln	Thr	Leu	Glu
	130				135							140				
	Thr	Leu	Asp	Lys	Val	Asp	Tyr	Ser	Lys	Tyr	Val	Ala	Val	Asn	Leu	Ala
	145				150						155					160
	Thr	Ser	His	Pro	His	Tyr	Asp	Ser	Ala	Gly	Asn	Ile	Leu	Asn	Met	Gly
				165						170					175	
30	Thr	Ser	Ile	Val	Asp	Lys	Gly	Arg	Thr	Lys	Tyr	Val	Leu	Phe	Lys	Ile
				180					185					190		
	Pro	Ser	Ser	Val	Pro	Glu	Lys	Glu	Lys	Lys	Lys	Ser	Cys	Phe	Lys	His
		195						200					205			
	Leu	Glu	Val	Val	Cys	Ser	Ile	Pro	Ser	Arg	Ser	Leu	Leu	Gln	Pro	Ser
35		210					215					220				
	Tyr	Tyr	His	Ser	Phe	Gly	Ile	Thr	Glu	Asn	Tyr	Ile	Val	Phe	Ile	Glu
	225				230					235						240
	Gln	Pro	Phe	Lys	Leu	Asp	Ile	Val	Lys	Leu	Ala	Thr	Ala	Tyr	Ile	Arg
				245						250					255	
40	Gly	Val	Asn	Trp	Ala	Ser	Cys	Leu	Ser	Phe	His	Lys	Glu	Asp	Lys	Thr
			260						265					270		
	Trp	Phe	His	Phe	Val	Asp	Arg	Lys	Thr	Lys	Lys	Glu	Val	Ser	Thr	Lys
		275						280					285			
	Phe	Tyr	Thr	Asp	Ala	Leu	Val	Leu	Tyr	His	His	Ile	Asn	Ala	Tyr	Glu
	290						295					300				
45	Glu	Asp	Gly	His	Val	Val	Phe	Asp	Ile	Val	Ala	Tyr	Arg	Asp	Asn	Ser
	305				310						315					320
	Leu	Tyr	Asp	Met	Phe	Tyr	Leu	Lys	Lys	Leu	Asp	Lys	Asp	Phe	Glu	Val
				325						330					335	
	Asn	Asn	Lys	Leu	Thr	Ser	Ile	Pro	Thr	Cys	Lys	Arg	Phe	Val	Val	Pro
50				340					345					350		
	Leu	Gln	Tyr	Asp	Lys	Asp	Ala	Glu	Val	Gly	Ser	Asn	Leu	Val	Lys	Leu
			355					360					365			

55

EP 1 031 627 A1

	Pro	Thr	Ser	Ala	Thr	Ala	Val	Lys	Glu	Lys	Asp	Gly	Ser	Ile	Tyr	Cys
	370						375					380				
5	Gln	Pro	Glu	Ile	Leu	Cys	Glu	Gly	Ile	Glu	Leu	Pro	Arg	Val	Asn	Tyr
	385					390					395					400
	Asp	Tyr	Asn	Gly	Lys	Lys	Tyr	Lys	Tyr	Val	Tyr	Ala	Thr	Glu	Val	Gln
					405					410					415	
	Trp	Ser	Pro	Val	Pro	Thr	Lys	Ile	Ala	Lys	Leu	Asn	Val	Gln	Thr	Lys
				420					425					430		
10	Glu	Val	Leu	His	Trp	Gly	Glu	Asp	His	Cys	Trp	Pro	Ser	Glu	Pro	Ile
		435						440					445			
	Phe	Val	Pro	Ser	Pro	Asp	Ala	Arg	Glu	Glu	Asp	Glu	Gly	Val	Val	Leu
	450					455					460					
	Thr	Cys	Val	Val	Val	Ser	Glu	Pro	Asn	Lys	Ala	Pro	Phe	Leu	Leu	Ile
15	465					470					475					480
	Leu	Asp	Ala	Lys	Thr	Phe	Lys	Glu	Leu	Gly	Arg	Ala	Thr	Val	Asn	Val
					485					490						
	Glu	Met	His	Leu	Asp	Leu	His	Gly	Met	Phe					495	
				500					505							
20																
25																
30																
35																
40																
45																
50																
55																

EP 1 031 627 A1

<210> 5
 <211> 529
 <212> PRT
 <213> Bos taurus

<400> 5

10	Glu	Glu	Leu	Ser	Ser	Pro	Leu	Thr	Ala	His	Val	Thr	Gly	Arg	Ile	Pro
	1				5					10					15	
	Leu	Trp	Leu	Thr	Gly	Ser	Leu	Leu	Arg	Cys	Phe	Thr	Gly	Pro	Gly	Leu
			20						25					30		
	Phe	Glu	Val	Gly	Ser	Glu	Pro	Phe	Tyr	His	Leu	Phe	Asp	Gly	Gln	Ala
			35					40					45			
15	Leu	Leu	His	Lys	Phe	Asp	Phe	Lys	Glu	Gly	His	Val	Thr	Tyr	His	Arg
	50					55						60				
	Arg	Phe	Ile	Arg	Thr	Asp	Ala	Tyr	Val	Arg	Ala	Met	Thr	Glu	Lys	Arg
	65				70						75					80
	Ile	Val	Ile	Thr	Glu	Phe	Gly	Phe	Thr	Thr	Cys	Ala	Phe	Pro	Asp	Pro
20					85					90					95	
	Cys	Lys	Asn	Ile	Phe	Ser	Arg	Phe	Phe	Ser	Tyr	Phe	Arg	Gly	Val	Glu
			100					105						110		
	Val	Thr	Asp	Asn	Ala	Leu	Val	Asn	Val	Tyr	Pro	Val	Gly	Glu	Asp	Tyr
			115					120					125			
25	Tyr	Ala	Cys	Thr	Glu	Thr	Asn	Phe	Ile	Thr	Lys	Ile	Asn	Pro	Glu	Thr
	130						135					140				
	Leu	Glu	Thr	Ile	Phe	Thr	Lys	Gln	Val	Asp	Leu	Cys	Asn	Tyr	Val	Ser
	145				150					155						160
	Val	Asn	Gly	Ala	Thr	Ala	His	Pro	His	Ile	Glu	Asn	Asp	Gly	Thr	Val
				165						170					175	
30	Tyr	Asn	Ile	Gly	Asn	Cys	Phe	Gly	Lys	Asn	Phe	Ser	Ile	Ala	Tyr	Asn
			180					185						190		
	Ile	Val	Lys	Ile	Pro	Pro	Leu	Gln	Ala	Asp	Lys	Glu	Asp	Pro	Ile	Ser
			195				200						205			
	Lys	Phe	Thr	Ser	Glu	Ile	Val	Val	Gln	Phe	Pro	Cys	Ser	Asp	Arg	Phe
35		210					215					220				
	Lys	Pro	Ser	Tyr	Val	His	Ser	Phe	Gly	Leu	Thr	Pro	Asn	Tyr	Ile	Val
	225				230					235						240
	Phe	Val	Glu	Thr	Pro	Val	Lys	Ile	Asn	Leu	Phe	Lys	Phe	Leu	Ser	Ser
				245						250					255	
40	Trp	Ser	Leu	Trp	Gly	Ala	Asn	Tyr	Met	Asp	Cys	Phe	Glu	Ser	Phe	Thr
			260						265						270	
	Asn	Glu	Thr	Met	Gly	Val	Trp	Leu	His	Ile	Ala	Asp	Lys	Lys	Arg	Lys
			275					280					285			
	Lys	Tyr	Leu	Asn	Asn	Lys	Tyr	Arg	Thr	Ser	Pro	Phe	Asn	Leu	Phe	His
		290					295					300				
45	His	Ile	Asn	Thr	Tyr	Glu	Asp	Asn	Gly	Phe	Leu	Ile	Val	Asp	Leu	Cys
	305					310					315					320
	Cys	Trp	Lys	Gly	Phe	Glu	Phe	Val	Tyr	Asn	Tyr	Phe	Thr	Leu	Tyr	Leu
				325						330					335	
	Ala	Asn	Leu	Arg	Glu	Asn	Trp	Glu	Glu	Val	Lys	Lys	Asn	Ala	Arg	Lys
50				340					345					350		
	Ala	Pro	Gln	Pro	Glu	Val	Arg	Arg	Tyr	Val	Leu	Pro	Leu	Asn	Ile	Asp
			355					360					365			

55

EP 1 031 627 A1

	Lys	Ala	Asp	Thr	Gly	Lys	Asn	Leu	Val	Thr	Leu	Pro	Asn	Thr	Thr	Ala
		370					375					380				
5	Thr	Ala	Ile	Leu	Cys	Ser	Asp	Glu	Phe	Thr	Thr	Ile	Trp	Leu	Glu	Pro
	385					390					395					400
	Glu	Val	Leu	Phe	Ser	Gly	Pro	Arg	Gln	Ala	Phe	Glu	Phe	Pro	Gln	Ile
					405					410					415	
	Asn	Tyr	Gln	Lys	Tyr	Cys	Gly	Lys	Pro	Tyr	Thr	Tyr	Ala	Tyr	Gly	Leu
				420					425						430	
10	Gly	Leu	Asn	His	Phe	Val	Pro	Asp	Arg	Leu	Cys	Lys	Leu	Asn	Val	Lys
			435					440					445			
	Thr	Lys	Glu	Thr	Trp	Phe	Thr	Val	Trp	Gln	Glu	Pro	Asp	Ser	Tyr	Pro
		450					455					460				
	Ser	Glu	Pro	Ile	Phe	Val	Ser	His	Pro	Asp	Ala	Leu	Glu	Glu	Asp	Asp
15		465				470					475					480
	Gly	Val	Val	Leu	Ser	Val	Val	Val	Ser	Pro	Gly	Ala	Gly	Gln	Lys	Pro
					485					490					495	
	Ala	Tyr	Leu	Leu	Ile	Leu	Asn	Ala	Lys	Asp	Leu	Ser	Glu	Val	Ala	Arg
				500					505					510		
20	Ala	Glu	Phe	Thr	Val	Glu	Ile	Asn	Ile	Pro	Val	Thr	Phe	His	Gly	Leu
			515					520					525			
	Phe															

25

30

35

40

45

50

55

<210> 6
 <211> 18
 <212> PRT
 5 <213> Gallus gallus
 <400> 6

10 Asn Lys Glu Glu His Pro Glu Pro Ile Lys Ala Glu Val Gln Gly Gln
 1 5 10 15
 Leu Pro

15 <210> 7
 <211> 18
 <212> PRT
 <213> Gallus gallus
 <400> 7

20 Asn Lys Glu Glu His Pro Glu Pro Ile Lys Ala Glu Val Gln Gly Gln
 1 5 10 15
 Leu Pro

25 <210> 8
 <211> 20
 <212> DNA
 30 <213> artificial sequence
 <220>
 <221> modified_base
 <222> 18
 35 <223> a is Inosine; primer
 <400> 8

40 aacaargarg ascayccaga 20

45

50

55

5 <210> 9
 <211> 21
 <212> DNA
 <213> artificial sequence

 <220>
 10 <221> modified_base
 <222> 7, 13
 <223> a is Inosine; primer

 <400> 9

15 sagctgaccc tgaacytcsg c

21

20 <210> 10
 <211> 25
 <212> DNA
 <213> artificial sequence

 <220>
 25 <223> primer

 <400> 10

30 tctgaattcc ggagcccata aaagc

25

35 Claims

1. Protein having β,β -carotene 15,15'-dioxygenase activity comprising an amino acid sequence which is identical or homologous to sequence ID no. 1 whereby the degree of homology to sequence ID no. 1 is at least 60%.
- 40 2. Protein according to claim 1 wherein the degree of homology is at least 70%.
3. Protein according to claim 1 wherein the degree of homology is at least 80%.
4. Protein according to claim 1 wherein the degree of homology is at least 90%.
- 45 5. Protein according to any one of claims 1 to 4 wherein the amino acid sequence is derived from chicken.
6. Nucleic acid sequence coding for a protein according to any one of claims 1 to 5.
- 50 7. Nucleic acid sequence according to claim 6 which comprises at least a part of the sequence of sequence ID no. 2.
8. Nucleic acid sequence according to claim 7 wherein the oligonucleotide has a length of at least 20 bases.
9. Nucleic acid sequence according to any one of claims 6 to 8 wherein the nucleic acid is desoxyribonucleic acid.
- 55 10. Nucleic acid sequence according to any one of claims 6 to 8 wherein the nucleic acid is an antisense ribonucleic acid.

11. Primer for the specific amplification of a gene coding for a protein having β,β -carotene 15,15'-dioxygenase activity which comprises a nucleic acid sequence according to any one of claims 6 to 9.

5 12. Probe for the detection of a gene coding for a protein having β,β -carotene 15,15'-dioxygenase activity which comprises a nucleic acid sequence according to any one of claims 6 to 9.

13. Testkit for the amplification and/or detection of a gene or a part thereof coding for β,β -carotene 15,15'-dioxygenase comprising at least one primer according to claim 11 and/or at least one probe according to claim 12.

10 14. Antibody characterized in that said antibody specifically reacts with a protein according to any one of claims 1 to 5.

15. Immunoassay for the detection and/or quantification of β,β -carotene 15,15'-dioxygenase which comprises at least one antibody according to claim 14.

15 16. Process for the production of vitamin A wherein β -carotene is enzymatically cleaved by a protein according to any one of claims 1 to 5.

17. Method for the introduction of the β,β -carotene 15,15'-dioxygenase cDNA into a host cell wherein a cDNA coding for a protein according to any one of claims 1 to 5 is inserted into a vector suitable for the host cell and said vector is introduced into the host cell.

18. Method according to claim 17 wherein the host cell is a plant cell.

19. Method according to claim 17 wherein the host cell is a prokaryotic cell.

20. Method according to claim 17 wherein the host cell is a yeast cell or another fungus.

21. Method according to claim 17 wherein the host cell is an alga.


22. Method according to claim 17 wherein the host cell is a mammalian cell.

23. Method according to claim 22 wherein the mammalian cell is a human cell.

24. Host cell obtainable by a method according to any one of claims 17 to 23.

25. Host cell according to claim 24 which comprises a β,β -carotene 15,15'-dioxygenase cDNA obtained from another species.

Figure 1






Figure 2

Figure 3
No. 2

Seq. ID

1 CGGATCCACT AGTAACGGCC GCCAGTGTGG TGGAATCCAT
CCTTCTATGT

51 AACAGGAAAG AGCTGTTCTT AGCCCAGAGA GGAGGGCACC
GTACGCCTGC

101 AGGAGCAGCT GGGTAGAGGA CACAGGAGAG CGATGGAGAC
AATATTTAAC

151 AGAAACAAAG AAGAGCATCC AGAGCCCATA AAAGCTGAGG
TGCAAGGTCA

201 GTTGCCCACT TGGTTGCAAG GGGTACTTCT CCGAAATGGC
CCAGGGATGC

251 ACACAATAGG GGACACTAAA TACAACCACT GGTTTGATGG
CTTGGCTCTG

301 CTGCACAGCT TCACGTTTAA AAATGGTGAA GTTTACTACA
GAAGTAAGTA

351 CCTCCGAAGT GACACATACA ACTGCAATAT AGAAGCAAAC
CGAATCGTGG

401 TGTCTGAGTT TGGAACCATG GCTTATCCGG ATCCATGCAA
AAACATATTT

451 GCCAAGGCAT TCTCATACTT ATCTCACACC ATTCCTGAGT
TCACGGACAA

501 CTGCCTGATC AACATTATGA AAAGTGGGGA TGATTATTAT
GCTACCAGTG

551 AGACTAACTT CATCAGAAAA ATTGATCCAC AGACTCTGGA
GACACTAGAT

601 AAGGTAGACT ACAGCAAATA TGTAGCTGTA AACTTGGCAA
CTTCTCACCC

651 ACACTATGAC AGTGCTGGAA ATATTCTCAA CATGGGTACT
TCAATTGTTG

701 ATAAAGGGAG AACAAAATAT GTTCTCTTTA AGATCCCTTC
CTCTGTACCA

751 GAAAAAGAAA AGAAGAAATC TTGTTTTTAAA CACCTGGAAG
TAGTATGCTC

801 CATCCCTTCT CGCTCCCTGC TCCAACCAAG CTACTACCAC
AGCTTTGGAA

851 TCACAGAAAA TTATATTGTG TTCATAGAGC AGCCATTTAA
ACTGGATATT

901 GTCAAACCTGG CAACTGCCTA CATCCGAGGT GTGAACTGGG
 CTTCCCTGCCT
 951 TTCCTTTTCAT AAGGAGGATA AGACGTGGTT TCACTTTGTA
 GACAGAAAGA
 1001 CGAAAAAAGA AGTATCCACC AAGTTTTACA CTGATGCTTT
 GGTGCTTTAT
 1051 CACCACATAA ATGCTTACGA AGAAGATGGC CACGTTGTTT
 TTGATATCGT
 1101 TGCCTACAGA GACAATAGCT TGTACGATAT GTTTTACTTA
 AAAAACTGG
 1151 ACAAAGACTT TGAAGTGAAC AACAAGCTTA CCTCCATCCC
 AACCTGCAAG
 1201 CGCTTTGTTG TGCCTCTGCA GTATGACAAG GATGCAGAAG
 TAGGTTCTAA
 1251 TTTAGTCAAA CTTCCAACCT CCGCAACTGC TGTAAGAGAA
 AAAGATGGCA
 1301 GCATCTATTG TCAACCTGAA ATATTATGTG AAGGGATAGA
 ACTGCCTCGT
 1351 GTCAAATATG ACTACAATGG CAAAAAATAC AAGTATGTCT
 ATGCAACAGA
 1401 AGTCCAGTGG AGCCCAGTTC CTACAAAGAT TGCAAAACTG
 AATGTCCAAA
 1451 CAAAGGAAGT ACTGCACTGG GGAGAAGACC ACTGCTGGCC
 CTCAGAGCCC
 1501 ATCTTTGTTC CCAGCCCCGA TGCAAGAGAA GAGGATGAAG
 GTGTTGTTTT
 1551 GACCTGTGTT GTGGTGTCTG AGCCAAATAA AGCACCCTTC
 CTACTCATCT
 1601 TGGATGCTAA AACATTCAAA GAATTGGGCC GAGCCACAGT
 TAACGTAGAA
 1651 ATGCATCTGG ACCTGCATGG GATGTTTATA CCACAGAATG
 ATTTGGGGGC
 1701 TGAGACGGAA TAAACGCTA TTGATCCGAC TACACAACT
 GAGACAACCT
 1751 TCTACTGAAC ATGAGTTAAT ATCCCTTTTA CCATTCAAGA
 ACAACCATAT
 1801 AACGACACAA AATGACTATG TATAATCTCT TAAATAATAG
 ATATAATCCT
 1851 TTTAAGGCAC AGCGATGAGT TTTACTACAG GTAACGATAT
 GCACAACCTGG

1901 CATATAACTA TTCCAAAAGA AGAAGAACGA TCAGTGTTTT
AGAAGTGCTA

1951 ATGTTGTACA TAACGGCGGC AGAGGGAACA GGAGAGAAAG
GTAACGGGAA

2001 TATTTAATAG AATATAGATT TCTGAGCAAA TGAAGTGCAG
TATTTATGGT

2051 GTGATGCATG GCATGAGTCA CATAGGTCTG CAGCTCATGT
ATCTTTTAGA

2101 GATCGTTTCA AGATTGCAGC TTGTGATGCA AGTTTTCTCC
AGCCAGAAAA

2151 CCTCATTTTA AACCATCTGC TACTGGTAAT TCATACCAAT
GCATTTTCTT

2201 GGTGCTCGAT TTACACTATA ACCAAAGTTA AGTATTACAT
TCAGGTGCTA

2251 CAACTTTCTA ATTTACAACC GAAACAAACA AGCAAACAGC
ACTTGCTTTG

2301 CTAATAACCC CATGGTGTAT TTTTCCTTTT TATGATGACA
AAACCAAGTA

2351 CATATGGTTT TATGTAGCAT TCAATTATAC TTCAGTGCTA
TTCCATCCTA

2401 ATGTTATAAG CAATTTGTAT TTAAATCAGT TTTCTTGAG
AATATCTGAC

2451 ATAACATTTT GTGTAATGAG ATGACTATGT TGTCTAAAGA
TGAACAGGAA

2501 TGTATCTTTT ATTAGTATTG TTAATTGTGT TACTAATACT
ATGCATATGA

2551 ATGAGAGCAA TGTATTTCTA GGAGAACTCA GATATACATT
CAACAATTTT

2601 TGTAGGTGAA AATGCATTTA CTGATGAAAG TTGAATCGTT
AATGAGGGAG

2651 AAAACTGGGT ATCCATCCAT CCAACTATGT TAGGTGTTCA
CCTGGTCTGT

2701 ATGTGACACC ACGCTGTTTG GGTATCTCTC ACTTTCACAT
ACCTGTTCTC

2751 ATGGTTTCTG CTACTCACTG TATTTTGCAG GAGAGAAACA
AAATGAAATC

2801 ACTGTCACTT ACTATCGCCC CATCACATAA GAACAATGGG
GCTTTGGTGA

2851 CTTGTTCATG ATTACATAAG ATGTTTGCAG CAGAGCAGCA
ATAGAACCAA

2901 CACCATCCAC AGTTCTTGCT TGCTCTGTGA TGAATCCCTT
TGCTGTCTTT

2951 ATGGTTTGCA TGTATGAAGA ATACACTGCC TAATTCTAAT
GTTAAAAAGT

3001 CACTGGGGTC AGATCTAGAG CTTAAGTAAG CAGTCTGGGG
TTTTCAAATG

3051 TTTATATGTT CCATAAAATG GAAATAAACA CCTCCATAAT
AAAAAAAAAA

3101 AAAAAAAAAA A

Figure 4
No. 1

Seq. ID

1 METIFNRNKE EHPEPIKAEV QGQLPTWLQG VLLRNGPGMH
TIGDTKYNHW

51 FDGLALLHSF TFKNGEVYYR SKYLRSDTYN CNIEANRIVV
SEFGTMAYPD

101 PCKNIFAKAF SYLSHTIPEF TDNCLINIMK TGDDYYATSE
TNFIRKIDPQ

151 TLETLDKVDY SKYVAVNLAT SHPHYDSAGN ILNMGTSIVD
KGRTKYVLFK

201 IPSSVPEKEK KKSCFKHLEV VCSIPSRSLI QPSYYHSFGI
TENYIVFIEQ

251 PFKLDIVKLA TAYIRGVNWA SCLSPHKEDK TWFFHFVDRKT
KKEVSTKFYT

301 DALVLYHHIN AYEEDGHVVF DIVAYRDNSL YDMFYLLKKLD
KDFEVNNKLT

351 SIPTCKRFVV PLQYDKDAEV GSNLVKLPTS ATAVKEKDGS
IYCQPEILCE

401 GIELPRVNYD YNGKKYKYVY ATEVQWSPVP TKIAKLNVT
KEVLHWGEDH

451 CWPSEPIFVP SPDAREEDED VVLTCVVVSE PNKAPFLIL
DAKTFKELGR

501 ATVNVEMHLD LHGMFIPQND LGAETE

Figure 5

Seq ID No. 4 and Seq ID No. 5

10 EEHPEPIKAÆVQGQLPTWLQGVLLR..NGPGMHTIGDTKYNHWF DGLALL
 57 || | : | | | : | | | | | | : | : | | | | |
 20 EELSSPLTAHV TGRIP LWTG SLLRCFTG PGLFEVGSEPFYHLFDGQALL
 69
 58 HSFTFKNGEVYYRSKYLRSDTYNCNIEANRIVVSEFG..TMAYPD PCKNI
 105 | | | | | | : : | | | | | | : | : | | | | |
 70 HKFDFKEGHV TYHRRFIRTDAYVRAMTEKRIVITEFGFTTCAFPDPCKNI
 119
 106 FAKAFSYLSHTIPEFTDNCLINIMKTGDDYATSETNFIRKIDPQTLET
 155 | : | | | | | | : : | : | | | | | | : | : | | | :
 120 FSRFFSYFRGV..EVT DNALVNVPVGEDYACTETNFITKINPETLETI
 167
 156 ..DKVDYSKYVAVN LATSHPHYDSAGNILNMGTSIVDKGR TKYVLFKIPS
 203 . | | | | | | | : . | : | . | : | | |
 168 FTKQVDLCNYVSVNGATAHPHIENDGTVYNIGNCFGKNFSIAYNIVKIPP
 217
 204 SVPEKEKKKSCFKHLEVVC SIPSRLLQPSYYHSFGITENYIVFIEQPFK
 253 : | | | | | | : | | | | | | : | | | | | | : | | |
 218 LQADKEDPISKFTS.EIVVQFPCSDRFKPSYVHSFGLTPNYIVFVETPVK
 266
 254 LDIVKLATAY.IRGVNWASCL.SFHKEDK.TWFHFVDRKTKKEVSTKFYT
 300 : : | : : : | : | | | | | | : | | | | : |
 267 INLFKFLSSWSLWGANYMDCFESFTNETMGVWLHIADKKRKKYLNNKYRT
 316
 301 DALVLYHHINAYEEDGHVVF DIVAYRDNSL...YDMFY LKKLDKDFE...
 344 | : | | | | | : . | : : : | | | | | | : | : |
 317 SPFNLFHHINTYEDNGFLIVDLCCWKGFEFVYNYFTLYLANLRENWEEVK
 366
 345 VNNKLTSIPTCKRFV VPLQYDKDAEVGSNLVKLP.TSATAV..KEKDGS
 391 | : | : | : | | | | : | | | | | | : | : |
 367 KNARKAPQPEVRRYVLEPLNIDK.ADTGKNLVTLPNTTATAILCSDEFTTI
 415
 392 YCQPEILCEG....IELPRVNYD.YNGKKYKYVYATEVQWSPVPTKIAKL
 436 : : | : | | | : | | | | | | | | : : |
 416 WLEPEVLFSGPRQAFEFQIN YQKYCGKPYTYAYGLGLNHF.VPDR LCKL
 464

484 437 NVQTKVLH..WGEDHCWPSEPIFVSPDAREEDEGVVLTCCVVVSEPNKA
 465 .|.||| | | :||| ||| ||| :||| . ||
 514 NVKTKETWFTVWQEPDSYPSEPIFVSHPDAL EDDGVVLSVVVSPGAGQK
 485 P.FLLILDAKTFKELGRA..TVNVEMHLDLHGMF 515
 | :|||.|| |. || || : . . ||:|
 515 PAYLLILNAKDLSEVARAEFTVEINIPVTFHGLF 548

BEST AVAILABLE COPY




Fig. 6 shows a 10% polyacrylamide gel with E.coli expressed β,β -carotene 15,15'-dioxygenase after affinity tag purification; lane 1 and lane 2: 2 fractions from the Co^{2+} -chelate column showing the main band at 60 kD; lane 3: low range molecular weight marker (Bio Rad).




Fig. 7 shows an HPLC profile of the reaction mixture at the end of an activity assay for the β,β -carotene 15,15'-dioxygenase following the procedure in example 1. The first peak in the chromatogram represents the internal standard, while the second peak corresponds to retinal as the only product formed during the central cleavage with β -carotene as substrate.




Fig. 8 confirms that the product peak in Fig. 7 is indeed retinal. A sample which was positive in the activity assay (green chromatogram) was spiked with retinal and analysed in second HPLC run (red chromatogram). The chromatograms of the two runs were then overlayed.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 00 10 2289

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
D, A	<p>DEVERY J ET AL: "Beta-carotene-15,15'-dioxygenase (EC 1.13.11.21) isolation reaction method and an improved assay procedure." BRITISH JOURNAL OF NUTRITION., vol. 72, no. 3, September 1994 (1994-09), pages 397-414, XP000914668 CAMBRIDGE., GB * abstract *</p>	1	<p>C12N15/53 C12N9/02 C12N1/21 C12N5/10 C12P23/00 C07K16/40 C12Q1/68 G01N33/573 /(C12N1/21, C12R1:19)</p>
A	<p>WO 93 23538 A (LUDWIG INST CANCER RES ;BAVIK CLAES OLOF (SE); ERIKSSON ULF (SE);) 25 November 1993 (1993-11-25) * page 21 - page 24 *</p>	1	<p>TECHNICAL FIELDS SEARCHED (Int.Cl.7)</p> <p>C12N C12P C07K G01N C12Q</p>
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 13 June 2000	Examiner Lejeune, R
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.82 (PM/C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 10 2289

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

13-06-2000

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9323538 A	25-11-1993	AU 4375493 A	13-12-1993
		US 5573939 A	12-11-1996
		US 5679772 A	21-10-1997
<hr/>			

EPO FORM PV459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

THIS PAGE BLANK (USPTO)